



Tillable Soils Map



State: **Iowa**
 County: **Butler**
 Location: **30-92N-17W**
 Township: **West Point**
 Acres: **148.15**
 Date: **1/22/2020**



Soils data provided by USDA and NRCS.

Area Symbol: IA023. Soil Area Version: 25									
Code	Soil Description	Acres	Percent of field	CSR2 Legend	Non-Irr Class *c	Irr Class *c	CSR2**	CSR	*n NCCPI Overall
377B	Dinsdale silty clay loam, 2 to 5 percent slopes	28.51	19.2%			Ile	94	90	88
175B	Dickinson fine sandy loam, 2 to 5 percent slopes	24.35	16.4%			IIle	50	55	60
173B	Hoopston fine sandy loam, 2 to 5 percent slopes	16.60	11.2%			Ile	55	55	68
41C	Sparta loamy fine sand, 5 to 9 percent slopes	10.83	7.3%			IVs Ile	34	25	50
173	Hoopston fine sandy loam, 0 to 2 percent slopes	10.45	7.1%			IIs	59	60	69
175C	Dickinson fine sandy loam, 5 to 9 percent slopes	9.70	6.5%			IIIle	45	40	61
84	Clyde silty clay loam, 0 to 3 percent slopes	7.68	5.2%			IIw	88	75	90
391B	Clyde-Floyd complex, 1 to 4 percent slopes	6.70	4.5%			IIw	87	72	88
682	Maxfield silt loam, 0 to 2 percent slopes	6.29	4.2%			IIw	83		73
41B	Sparta loamy fine sand, 2 to 5 percent slopes	6.25	4.2%			IVs Ile	39	40	50
184	Klinger silty clay loam, 1 to 4 percent slopes	5.64	3.8%			Iw	95	95	88
83C2	Kenyon loam, 5 to 9 percent slopes, eroded	5.57	3.8%			IIIle	84	68	68
150B	Hanska loam, 1 to 4 percent slopes	5.36	3.6%			IIw	49	59	79
394C2	Ostrander loam, 5 to 9 percent slopes, eroded	2.28	1.5%			IIIle	73	68	58
41D	Sparta loamy fine sand, 9 to 14 percent slopes	1.57	1.1%			VIIs	8	15	48
174B	Bolan loam, 2 to 5 percent slopes	0.22	0.1%			IIs	64	70	86
118	Garwin silty clay loam, 0 to 2 percent slopes	0.15	0.1%			IIw	90	95	95
Weighted Average							65.7	*	*n 71.2

**IA has updated the CSR values for each county to CSR2.

*- CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

*n: The aggregation method is "Weighted Average using major components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.