

Soils Map





State: County: Lafayette 2-2N-3E Location: Township: Darlington

Acres: 118.9 8/30/2021 Date:







Soils data provided by USDA and NRCS.

| Area Symbol: WI065, Soil Area Version: 17 | | | | | | |
|---|--|-------|------------------|----------------------|------------------|------------------|
| Code | Soil Description | Acres | Percent of field | NCCPI Overall Legend | Non-Irr Class *c | *n NCCPI Overall |
| PaC2 | Palsgrove silt loam, 6 to 12 percent slopes, moderately eroded | 40.08 | 33.7% | | IIIe | 74 |
| FdB2 | Fayette silt loam, 2 to 6 percent slopes, moderately eroded | 35.85 | 30.2% | | lle | 86 |
| PaB2 | Palsgrove silt loam, 2 to 6 percent slopes, moderately eroded | 9.35 | 7.9% | | lle | 76 |
| FdC2 | Fayette silt loam, 6 to 12 percent slopes, moderately eroded | 8.43 | 7.1% | | IIIe | 83 |
| PaD2 | Palsgrove silt loam, 12 to 20 percent slopes, moderately eroded | 6.63 | 5.6% | | IVe | 68 |
| HvA | Huntsville silt loam, 0 to 2 percent slopes | 5.17 | 4.3% | | llw | 80 |
| ChB | Chaseburg silt loam, moderately well drained, 2 to 6 percent slopes | 4.34 | 3.7% | | llw | 85 |
| DsD2 | Newglarus silt loam, moderately deep, 12 to 20 percent slopes, moderately eroded | 4.32 | 3.6% | | IVe | 51 |
| An | Arenzville silt loam, 0 to 3 percent slopes, occasionally flooded | 1.94 | 1.6% | | llw | 88 |
| DsC2 | Newglarus silt loam, moderately deep, 6 to 12 percent slopes, moderately eroded | 1.19 | 1.0% | | IIIe | 56 |
| DsB2 | Newglarus silt loam, moderately deep, 2 to 6 percent slopes, moderately eroded | 0.68 | 0.6% | | lle | 57 |
| La | Lawson silt loam | 0.68 | 0.6% | | IIIw | 60 |
| On | Orion silt loam, 0 to 3 percent slopes, occasionally flooded | 0.24 | 0.2% | | llw | 90 |
| Weighted Average | | | | | | *n 77.8 |

^{*}n: The aggregation method is "Weighted Average using all components" *c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.