



Total Soils Map



Soils data provided by USDA and NRCS.



State: **Wisconsin**
 County: **Lafayette**
 Location: **35-4N-5E**
 Township: **Blanchard**
 Acres: **420.43**
 Date: **5/1/2022**



Maps Provided By:



Area Symbol: WI065, Soil Area Version: 18

Code	Soil Description	Acres	Percent of field	NCCPI Overall Legend	Non-Irr Class *c	*n NCCPI Overall
MfC2	Mifflin soils, 6 to 12 percent slopes, moderately eroded	49.29	11.7%		IIIe	68
NfD2	Northfield loam, 12 to 20 percent slopes, moderately eroded	44.05	10.5%		VIe	49
NfE2	Northfield loam, 20 to 30 percent slopes, moderately eroded	28.21	6.7%		VIIe	12
DsC2	Newglarus silt loam, moderately deep, 6 to 12 percent slopes, moderately eroded	25.22	6.0%		IIIe	54
DsB2	Newglarus silt loam, moderately deep, 2 to 6 percent slopes, moderately eroded	25.19	6.0%		Ile	56
Ba	Boaz silt loam	23.65	5.6%		IIIw	87
DsD2	Newglarus silt loam, moderately deep, 12 to 20 percent slopes, moderately eroded	23.38	5.6%		IVe	50
ChB	Chaseburg silt loam, moderately well drained, 2 to 6 percent slopes	20.16	4.8%		IIw	85
NoC2	Northfield sandy loam, 6 to 12 percent slopes, moderately eroded	18.16	4.3%		IVe	43
NfC2	Northfield loam, 6 to 12 percent slopes, moderately eroded	17.72	4.2%		IVe	53
An	Arenzville silt loam, 0 to 3 percent slopes, occasionally flooded	15.10	3.6%		IIw	88
PaC2	Palsgrove silt loam, 6 to 12 percent slopes, moderately eroded	14.97	3.6%		IIIe	74
GoB2	Gotham loamy fine sand, 2 to 6 percent slopes, moderately eroded	14.16	3.4%		IVs	50
MfD2	Mifflin soils, 12 to 20 percent slopes, moderately eroded	11.64	2.8%		IVe	62
Sb	Sable silt loam, benches	10.18	2.4%		IIw	68
NoD2	Northfield sandy loam, 12 to 20 percent slopes, moderately eroded	9.58	2.3%		VIe	40
1195E	Elk mound-Northfield complex, 20 to 30 percent slopes, very rocky	8.76	2.1%		VIIe	10
On	Orion silt loam, 0 to 3 percent slopes, occasionally flooded	8.35	2.0%		IIw	90
NoE2	Northfield sandy loam, 20 to 30 percent slopes, moderately eroded	8.02	1.9%		VIIe	10
FdB2	Fayette silt loam, 2 to 6 percent slopes, moderately eroded	6.00	1.4%		Ile	86
1195F	Elk mound-Northfield complex, 30 to 60 percent slopes, very rocky	5.30	1.3%		VIIe	10
MeA	Meridian sandy loam, 0 to 2 percent slopes	5.17	1.2%		IIIIs	67
DbB	Dakota loam, mottled subsoil variant, 0 to 3 percent slopes	4.61	1.1%		IIw	70
W	Water	4.54	1.1%			
HmC2	Hixton loam, 6 to 12 percent slopes, moderately eroded	4.41	1.0%		IIIe	59

HmD2	Hixton loam, 12 to 20 percent slopes, moderately eroded	3.85	0.9%		IVe	54
BoD2	Boone fine sand, 6 to 20 percent slopes, moderately eroded	2.81	0.7%		VIIIs	18
NoB2	Northfield sandy loam, 2 to 6 percent slopes, moderately eroded	2.03	0.5%		IIIs	44
HvA	Huntsville silt loam, 0 to 2 percent slopes	1.69	0.4%		IIw	85
Ma	Marsh	1.40	0.3%		VIIw	
MfB2	Mifflin soils, 2 to 6 percent slopes, moderately eroded	1.29	0.3%		IIe	68
La	Lawson silt loam	0.67	0.2%		IIIw	62
FeC2	Fayette silt loam, valleys, 6 to 12 percent slopes, moderately eroded	0.66	0.2%		IIle	85
ChA	Chaseburg silt loam, moderately well drained, 0 to 2 percent slopes	0.21	0.0%		IIw	86
Weighted Average					3.86	*n 55.5

*n: The aggregation method is "Weighted Average using all components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.