

Tillable Soils Map





State: lowa
County: Greene
Location: 36-84N-29W
Township: Junction
Acres: 152.02

Date:



7/3/2023





Soils data provided by USDA and NRCS.

Area Symbol: IA073, Soil Area Version: 28										
Code	Soil Description	Acres	Percent of field	CSR2 Legend	Non-Irr Class *c	*i Corn Bu	*i Soybeans Bu	CSR2**	CSR	*n NCCPI Overall
L107	Webster clay loam, Bemis moraine, 0 to 2 percent slopes	90.36	59.4%		llw	0.0	0.0	88		84
L55	Nicollet loam, 1 to 3 percent slopes	27.51	18.1%		le	0.0	0.0	91		84
6	Okoboji silty clay loam, 0 to 1 percent slopes	10.74	7.1%		IIIw	185.6	53.8	59	58	76
34B	Estherville sandy loam, 2 to 6 percent slopes	7.44	4.9%		IIIs	80.0	23.2	27	20	49
506	Wacousta silty clay loam, 0 to 1 percent slopes	4.83	3.2%		IIIw	198.4	57.5	74	77	87
L507	Canisteo clay loam, Bemis moraine, 0 to 2 percent slopes	3.53	2.3%		llw	0.0	0.0	87		81
108	Wadena loam, 0 to 2 percent slopes	2.04	1.3%		lls	80.0	23.2	56		62
L638C2	Clarion-Storden complex, Bemis moraine, 6 to 10 percent slopes, moderately eroded	1.98	1.3%		Ille	0.0	0.0	75		68
135	Coland clay loam, 0 to 2 percent slopes, occasionally flooded	1.93	1.3%		llw	198.4	57.5	76	81	83
L95	Harps clay loam, Bemis moraine, 0 to 2 percent slopes	1.66	1.1%		llw	0.0	0.0	75		80
Weighted Average					1.98	26.9	7.8	82.1	*-	*n 81.2

^{**}IA has updated the CSR values for each county to CSR2.

Soils data provided by USDA and NRCS.

^{*-} CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

^{*}i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

^{*}n: The aggregation method is "Weighted Average using all components"

^{*}c: Using Capabilities Class Dominant Condition Aggregation Method