

Tillable Soils Map





State: Iowa County: Greene

Location: 33-82N-32W

Township: Willow Acres: 78.99 Date: 6/23/2021







Soils data provided by USDA and NRCS.

| Area Symbol: IA073, Soil Area Version: 26 | | | | | | | | | | |
|---|---|-------|------------------|----------------|---------------------|------------|-------------|--------|-----|---------------------|
| Code | Soil Description | Acres | Percent of field | CSR2 Legend | Non-Irr Class *c | *i Corn | *i Soybeans | CSR2** | CSR | *n NCCPI Overall |
| L638C2 | Clarion-Storden complex, Bemis moraine, 6 to 10 percent slopes, moderately eroded | 35.00 | 44.3% | | Ille | 0 | 0 | 75 | | 70 |
| L138B2 | Clarion loam, Bemis moraine, 2 to 6 percent slopes, moderately eroded | 9.53 | 12.1% | | lle | 0 | 0 | 85 | | 72 |
| L138C2 | Clarion loam, Bemis moraine, 6 to 10 percent slopes, moderately eroded | 8.10 | 10.3% | | Ille | 0 | 0 | 83 | | 67 |
| L138B | Clarion loam, Bemis moraine, 2 to 6 percent slopes | 7.69 | 9.7% | | lle | 220.8 | 64 | 88 | | 80 |
| L55 | Nicollet loam, 1 to 3 percent slopes | 7.07 | 9.0% | | le | 0 | 0 | 91 | | 85 |
| 585B | Coland-Spillville complex, 1 to 5 percent slopes | 3.07 | 3.9% | | llw | 195.2 | 56.6 | 74 | 70 | 92 |
| L638D2 | Omsrud-Storden complex, Bemis moraine, 10 to 16 percent slopes, moderately eroded | 2.75 | 3.5% | | IVe | 0 | 0 | 53 | | 67 |
| 485B | Spillville loam, 2 to 5 percent slopes | 2.09 | 2.6% | | lle | 200 | 58 | 88 | 86 | 89 |
| L62D2 | Storden loam, Bemis moraine, 10 to 16 percent slopes, moderately eroded | 1.78 | 2.3% | | IVe | 0 | 0 | 41 | | 65 |
| L107 | Webster clay loam, Bemis moraine, 0 to 2 percent slopes | 1.23 | 1.6% | | llw | 0 | 0 | 88 | | 84 |
| L138C | Clarion loam, Bemis moraine, 6 to 10 percent slopes | 0.68 | 0.9% | | IIIe | 0 | 0 | 84 | | 77 |
| Weighted Average | | | | | | 34.4 | 10 | 78.8 | *- | *n 73.7 |

^{**}IA has updated the CSR values for each county to CSR2.

Soils data provided by USDA and NRCS.

^{*-} CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

^{*}i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

^{*}n: The aggregation method is "Weighted Average using all components"

^{*}c: Using Capabilities Class Dominant Condition Aggregation Method