



# Current Tillable Soils Map



State: **Iowa**  
 County: **Greene**  
 Location: **34-82N-30W**  
 Township: **Franklin**  
 Acres: **141.43**  
 Date: **9/13/2021**



Area Symbol: IA073, Soil Area Version: 26

Code	Soil Description	Acres	Percent of field	CSR2 Legend	Non-Irr Class *c	*i Corn	*i Soybeans	CSR2**	CSR	*n NCCPI Overall
L55	Nicollet loam, 1 to 3 percent slopes	37.90	26.8%		Ie	0	0	91		85
259	Biscay clay loam, 0 to 2 percent slopes	21.39	15.1%		IIw	80	23.2	52	78	69
L107	Webster clay loam, Bemis moraine, 0 to 2 percent slopes	19.96	14.1%		IIw	0	0	88		84
L138B	Clarion loam, Bemis moraine, 2 to 6 percent slopes	16.36	11.6%		Ile	220.8	64	88		80
L138C2	Clarion loam, Bemis moraine, 6 to 10 percent slopes, moderately eroded	15.90	11.2%		IIIe	0	0	83		67
L138B2	Clarion loam, Bemis moraine, 2 to 6 percent slopes, moderately eroded	11.10	7.8%		Ile	0	0	85		72
639D2	Salida-Storden complex, 9 to 14 percent slopes, moderately eroded	5.77	4.1%		IVe	80	23.2	26	10	48
L507	Canisteo clay loam, Bemis moraine, 0 to 2 percent slopes	5.18	3.7%		IIw	0	0	87		81
L638C2	Clarion-Storden complex, Bemis moraine, 6 to 10 percent slopes, moderately eroded	4.78	3.4%		IIIe	0	0	75		70
203	Cylinder loam, 0 to 2 percent slopes	3.09	2.2%		IIIs	80	23.2	58	79	73
<b>Weighted Average</b>						<b>42.7</b>	<b>12.4</b>	<b>78.9</b>	<b>*</b>	<b>*n 76.4</b>

\*\*IA has updated the CSR values for each county to CSR2.

\*. CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

\*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

\*n: The aggregation method is "Weighted Average using all components"

\*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.