

Current Tillable Soils Map





State: lowa
County: Grundy
Location: 33-88N-16W
Township: Lincoln
Acres: 172.02

Date:



6/7/2023





Soils data provided by USDA and NRCS.

Area Symbol: IA075, Soil Area Version: 28										
Code	Soil Description	Acres	Percent of field	CSR2 Legend	Non-Irr Class *c	*i Corn Bu	*i Soybeans Bu	CSR2**	CSR	*n NCCPI Overall
120B	Tama silty clay loam, 2 to 5 percent slopes	54.28	31.6%		lle	232.0	67.3	95	95	97
133	Colo silty clay loam, 0 to 2 percent slopes, occasionally flooded	41.02	23.8%		llw	204.8	59.4	78	80	77
11B	Colo-Ely complex, 0 to 5 percent slopes	19.82	11.5%		llw	80.0	23.2	86		92
377B	Dinsdale silty clay loam, 2 to 5 percent slopes	16.88	9.8%		lle	232.0	67.3	94	90	87
119B	Muscatine silty clay loam, 2 to 5 percent slopes	15.06	8.8%		lle	232.0	67.3	95	95	96
221	Klossner muck, 1 to 4 percent slopes	8.48	4.9%		IIIw	80.0	23.2	48	50	88
241C2	Saude-Lilah complex, 5 to 14 percent slopes, moderately eroded	5.23	3.0%		Ille	80.0	23.2	35	28	53
120C	Tama silty clay loam, 5 to 9 percent slopes	4.12	2.4%		Ille	216.0	62.6	90	80	93
933	Sawmill silty clay loam, shallow loess, 0 to 2 percent slopes, occasionally flooded	3.22	1.9%		llw	204.8	59.4	78		77
7	Wiota silt loam, 0 to 2 percent slopes, rarely flooded	1.97	1.1%		lw	233.6	67.7	100	95	93
241B	Saude-Lilah complex, 2 to 5 percent slopes	1.13	0.7%		lle	80.0	23.2	48	58	65
88	Nevin silty clay loam, 0 to 2 percent slopes	0.81	0.5%		lw	227.2	65.9	95	95	98
Weighted Average					2.09	194	56.3	85	*-	*n 88.1

^{**}IA has updated the CSR values for each county to CSR2.

^{*-} CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

^{*}i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

^{*}n: The aggregation method is "Weighted Average using all components"

^{*}c: Using Capabilities Class Dominant Condition Aggregation Method Soils data provided by USDA and NRCS.